

Section 1

*Lecturer: Larry Blume**TA: Abhi Ananth*

* These notes develop Fikri Pitsuwan's notes from 2017.

Logistics

- OH: Thus 4-6 pm
- Material available at: <https://abhiananthecon.github.io/teaching/>
- Same link for office hours and sections
- Please email me with subject header 6100 to be a part of the mailing list
- Thu 6pm deadline for topic suggestions
- Questions?

Today we will look at:

1. Farka's lemma
2. Canonical and standard form
3. Vertex theorem


1 Review

Let's start with Farka's lemma. It states that for any $A \in \mathbb{R}^{m \times n}$ and $b \in \mathbb{R}^m$, **exactly one** of the following **will hold**:

- There is some $x \in \mathbb{R}^n$ satisfying $x \geq 0$ and $Ax = b$.
- There is some $y \in \mathbb{R}^m$ satisfying $yA \geq 0$ and $yb < 0$.

Farkas's Lemma:

Suppose $A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix} \begin{matrix} \vec{a}_1 \\ \vec{a}_2 \end{matrix}$; $b = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$



If \vec{b} lies in purple area, $A\vec{x} = b, \vec{x} \geq \vec{0}$ has a solution. Eg $\vec{b} = (1, 1) \rightarrow x^*(\vec{b}) = (\frac{2}{5}, \frac{1}{5})$


Else, it has no solution. Eg $\vec{b} = (3, 1)$.

Farkas's lemma says that when $\{\vec{x} \geq \vec{0} : A\vec{x} = b\} = \emptyset$,

then

$$\{\vec{y} : \vec{y}A \geq \vec{0}, \vec{y}b < 0\} \neq \emptyset.$$

$$\tilde{a}_1 = [1 \ 2] ; \tilde{a}_2 = [2 \ 1]$$



Why it matters?

- Neat application of the separating hyperplane theorem
- Easy to verify criterion for feasibility of a linear program

A linear program can be written in *canonical form* as

$$\begin{aligned} v_p(b) &= \max c \cdot x \\ \text{s. t. } Ax &\leq b \\ x &\geq 0 \end{aligned}$$

where $c \in \mathbb{R}^n$, $x \in \mathbb{R}^n$, A is an $m \times n$ matrix, $b \in \mathbb{R}^m$. Any linear program can also be written in *standard form* as

$$\begin{aligned} v_p(b) &= \max c \cdot x \\ \text{s. t. } Ax &= b \\ x &\geq 0 \end{aligned}$$

Given an inequality constraint $2x_1 + 3x_2 \leq 5$ and $x_1, x_2 \geq 0$, we can introduce a slack variable $z_1 \geq 0$, so that the constraint becomes $2x_1 + 3x_2 + z_1 = 5$. Given an equality constraint $x_1 + 2x_2 = 3$, we can express this as $x_1 + 2x_2 \leq 3$ and $-x_1 - 2x_2 \leq -3$. A linear program with no non-negativity constraint can be dealt with by expressing $x = y - z$ with $y \geq 0$ and $z \geq 0$.

Here are some important definitions in linear programming.

Definition 1. Any $x \in \mathbb{R}^n$ is called a *solution*.

Definition 2. For a linear program in canonical form, $C = \{x \in \mathbb{R}^n : Ax \leq b, x \geq 0\}$ is called the *constraint set* or the *feasible set*. Any $x \in C$ is called a *feasible solution*.

Definition 3. A vector x that actually solves the linear program, i.e., $x \in C$ and $c \cdot x \geq c \cdot x'$ for all $x' \in C$ is called an *optimal solution*.

Definition 4. A vector $x \in C$ is a vertex of C if and only if there is no $y \neq 0$ such that $x + y$ and $x - y$ are both in C .

Theorem (Vertex Theorem). For a linear program in standard form with feasible solutions, a vertex exists and if $v_p(b) < \infty$ and $x \in C$, then there is a vertex x' such that $c \cdot x' \geq c \cdot x$.

Notes

A good reference on linear programming is *Introduction to Linear Optimization* by Bertsimas and Tsitsiklis.

2 Problems

Problem 1. Consider the following linear program

$$\begin{aligned} & \max 2x_1 + x_2 \\ & \text{s. t. } x_1 + x_2 \leq 1 \\ & \quad x_1 \geq 0, x_2 \geq 0 \end{aligned}$$

- (a) Express the linear program in canonical form and draw the constraint set and solve the problem graphically.
- (b) Express the linear program in standard form and draw the constraint set.
- (c) Verify that the vertex theorem applies. Use the vertex theorem to find an optimal solution of the linear program.

Problem 2. Consider the following linear program

$$\begin{aligned} & \max 2x_1 + x_2 \\ & \text{s. t. } x_1 + x_2 \leq 1 \\ & \quad 2x_2 - x_1 \geq -1 \end{aligned}$$

- (a) Draw the constraint set as given and solve the problem graphically.
- (b) Solve the problem using the Kuhn-Tucker formulation
- (c) Express the linear program in canonical form and in standard form.

Problem 3. Consider a utility maximization problem with $u(x) = \sum_{i=1}^n \alpha_i x_i$, where $\alpha_i > 0$ for all i .

- (a) Express the problem as a linear program in canonical form. What is the feasible set? What are c , A , and b ?
- (b) Solve the UMP for $n = 2$ using the Kuhn-Tucker formulation with $\alpha_1 = 3$, $\alpha_2 = 2$, $p_1 = 3$, $p_2 = 1$, $w = 3$. Verify your solution graphically.